

WORLD INTELLECTUAL PROPERTY ORGANIZATION



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7: G06F 17/60

(11) International Publication Number:

WO 00/17799

(43) International Publication Date:

30 March 2000 (30.03.00)

(21) International Application Number:

PCT/US99/22019

(22) International Filing Date: 22 September 1999 (22.09.99)

(30) Priority Data:

09/159,219

23 September 1998 (23.09.98)

(71) Applicant (for all designated States except US): HEALTH HERO NETWORK, INC. [US/US]; Suite 111, 2570 West El Camino Real, Mountain View, CA 94040 (US).

(72) Inventor; and


(75) Inventor/Applicant (for US only): BROWN, Stephen, J. [-/US]; 3324 Woodside Road, Woodside, CA 94062 (US).

(74) Agent: GRAHAM, Lawrence, D.; Black Lowe & Graham, PLLC, 816 Second Avenue, Scattle, WA 98104 (US).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AM, AL, BT, KU, KL, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FK, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of

(54) Title: DYNAMIC MODELING AND SCORING RISK ASSESSMENT



(57) Abstract

The invention provides for modeling and scoring risk-assessment and a set of insurance products derived therefrom. Risk indicators are determined at a selected time. A population is assessed at that time and afterward for those risk indicators and for consequences associated therewith. Population members are coupled to client devices for determining risk indicators and consequences. A server receives data from each client, and in response thereto and in conjunction with an expert operator, (1) teasesses weights assigned to the risk indicators, (2) determines new risk indicators, (3) determines new measures for determining risk indicators and consequences, and (4) presents treatment options to each population member. The server determines, in response to the data from each client, and possibly other data, a measure of risk for each indicated consequence or for a set of such consequences. The server provides this measure with regard to each population member, or with regard to population subsets. The expert operator uses this measure to determine either (1) an individual course of treatment, (2) a resource utilization review model, (3) a risk-assessment model, or (4) an insurance pricing model, for each individual population member or for selected population subsets. Information requested by the client, information determined and presented by the server, and responsive measurements, are adapted dynamically to changing population aspects or changing population membership, or of an external environment having relevance to the population.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

		ES	Spein .	LS	Lesotho	SI	Slovenia
	Albania	ñ	Finland	LT	Lithuania	SK	Slovakia
	Amenia	FR	France	ш	Lexembuarg	SN	Senegal
	Ausma	GA.	Gabon	LV	Latvia	SZ	Swaziland
	Australia	CB	United Kingdom	MC	Monaco	170	Chad
	Azerbaijan	GE.	Georgia	MD	Republic of Moldova	TC	Togo
	Bosnia and Herzegovina		Ghana .	MG	Madagascar .	, TJ	Tajikistan
	Barbados	GH	Guinea -	MK	The former Yugoslav	TM	Terkmenistan
	Belgium	CN	•		Republic of Macedonia	TR	Torkey
BF	Burkina Faso	GR	Greece	ML	Mali	17	Trinidad and Tobaco
BC	Bulgaria	HU	Hungary	MN	Mongolia	UA	Ukraine
BJ	Benin	IE	keland	MR	Mauritania	UG	Uganda
BR	Brazil	IL	Israel		Malawi	US	United States of Americ
BY	Belares	LS	keland	MW	Mexico	UZ.	Uzbekistan
CA	Canada	IT	kely .	MX.		VN	Vict Nam
	Central African Republic	JP	Japan	NÉ	Niger	YU	Yugoslavia
	Congo	KE	Kenya	NL	Netherlands	ZW	Zimbabwe
	Swirzerland	KG	Kyrgyzsian	NO .	Norway	ZW	Zanosowe
•	Core d'Ivoire	KP	Democratic People's	NZ.	New Zealand		
	Cameroon :		Republic of Korea	PL	Poland	· · · · ·	
	China	KR	Republic of Korea	PT	Portuga3		
	Cuba .	KZ.	Kazaksian	RO	Romania		
	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
	Creen repons	ш	Liechenstein	SD	Sudan		
	Denmark	LK	Sri Lanka	S€	Sweden		
	Esonia	LR	Liberia	· SG	Singapore		•
EE	P.SIOMIA						

	·	•	· ·	•			
I.				•			
			•				
					•		
2			•				
				• •			•
3					• •		
-	•			•			
				÷	. •	•• •	• .
4			•		· '		
		•	• • .	•			
5		•			•		
			•	•		•	
							•
0		•					
٠.							
7		•					•
		•					
8				•	•		٠.
U		•			•		
_			Title of t	he Invention		• •	
9		. ' '	110.01.2				
			•	•			.*
10			•		•		
11		Dynamic	Modeling and	I Scoring Ri	sk Assessme	nt	
•		•			.*	•	
,,	•			•		•	
12	•	•			•		
			Background	of the Inven	tion		
13			Dackground	Of the miven	<u>ation</u>		
							•
14							
15	1.	Field of the Invention	•				
							٠.
,,						·	
16							•
		This invention relates	to computer ex	retemic and d	lata structure	s for modelin	ne and
17		THIS INVENTION LEIGHES	to computer sy	stems and t	iota su uctul c	, tot modelii	-D
			•		•		
18	scorin	g risk assessment, such	as insurance r	isk.			
10					•		

PCT/US99/22019

2. Related Art

2

In the insurance industry and in other fields in which risk is assessed (including such diverse fields as medical treatment, financial modeling and portfolio management, and environmental impact regulation), it is known to develop and use a risk-assessment model of a population. The risk-assessment model provides a technique for determining which population members are more subject or less subject to particular risks (or to an aggregate of risks) than the norm for that population. For example, in life insurance underwriting, it is known to evaluate past and present medical data so as to determine what insurance premium the underwriter wishes to charge.

11

12

13

14

15

16

While these known methods generally achieve the goal of assessing risk for particular individuals in comparison to a population norm, they have the drawback of making a risk assessment that is fixed at a particular point in time. That is, these risk-assessment models rely on static data, in particular (1) static data about the individual population member, (2) static data about the population norm, and (3) static data about risks associated or correlated with the data about the individual population member. However, risk for individual population members depends not only on their present data, but also on their future data, including both data about behavior and environment.

20

19

A first type of problem for the known art includes those individuals that have a progressive disease or degenerative condition, in which the disease or condition

progresses at a rate that is responsive to behavior or environment of the individual. For such individuals, risk is more accurately evaluated as a function of behavior measured over time and environment measured over time, rather than as a static value that is a function only of present behavior and environment. For example, a first patient with diabetes can proceed with relatively small risk if that first patient is aware of and active in management of behavioral and environmental risk factors. In contrast, an otherwise identical second patient will have significantly greater risk if that second patient is either unaware of, or unable or unwilling to take charge of, behavioral and environmental risk factors.

10.

Related to this first type of problem is the problem of determining trends for individual risk-assessment. For example, an individual with a history of diabetes may suffer a significant increase or decrease in effects thereof, due at least in part to that patient's actions with regard to behavioral and environmental risk factors. Similarly to the first type of problem, that individual will be rationally assessed a significantly greater or lesser risk than originally, if the new facts were known to the underwriter. Such trends may differ significantly from any trends that might have been discerned from past medical history alone; such trends may also themselves involve genetic, environmental, or behavioral components, or some combination thereof.

A second type of problem for the known art includes individuals whose riskassessment significantly changes due to the vicissitudes of their life trajectory. This can WO 00/17799 PCT/US99/22019

include progression of a disease or condition, responsive at least in part to behavioral or environmental factors. For a more striking example, an individual may suffer a myocardial infarction, or become infected with an HIV variant. Similarly to the first type of problem, that individual would be rationally assessed a significantly greater risk than originally, if the new facts were known to the underwriter. Alternatively, an individual may be successfully treated for a "curable" disease such as Hodgkin's disease or some forms of cancer. Such vicissitudes of life trajectory may themselves involve genetic, environmental, or behavioral components, or some combination thereof.

A third type of problem for the known art includes individuals who significantly change their behavior or environment, particularly when those individuals are susceptible to the elements of their behavior or environment they change. For example, an individual with diabetes can determine to alter their diet favorably or unfavorably. For a more striking example, an individual may take up smoking or skydiving as habits. That individual will become a significantly greater risk than the underwriter originally assessed.

Moreover, new medical research may indicate risk factors that were not known at the time risk for the individual was originally assessed. These could include past medical information not known at the time to be important, tests available in the future for risk factors not known at the time at all, or changes in the medical history of the individual that place that individual in different risk factor categories. Such past medical

information or risk factors may themselves involve genetic, environmental, or behavioral elements, or some combination thereof.

3

Accordingly, it would be advantageous to collect feedback from individual

population members, whether on a periodic or aperiodic basis, and whether prompted by

selected events or not. Such feedback would allow underwriters or other risk-assessment

or risk-management personnel to determine specific risk-related information about each

individual population member, and to adjust (such as to make more accurate or precise)

insurance models and risk-assessment models to fit the new data. Such feedback enables

the advantage of providing information about the time-varying nature of individual

measures which can be used in the dynamic risk assessment model presented in the

present invention. For instance, a weight gain of 10 pounds per year, an increase in

diastolic blood pressure of 10 points per year, and a increase of cholesterol of 10 points

per year could be tracked over time and would yield health risk information.

15

16

20

To achieve this advantage, a first aspect of the invention is that feedback is collected by a client-server system in which data is requested or required from population members. A server device, responsive to a risk-assessment model, prompts a client device supplied to population members to request information from population members, in order to determine whether aggregate measures or individual measures of risk-assessment remain in coherence with the model. The client device collects the data and supplies it to the server device, which can, in response to dynamically collected data,

adjust the model, adjust risk assessments for selected population members (or groups thereof), or determine further information to collect from population members.

Upon achieving this advantage, a second aspect of the invention is to provide a set of superior risk-assessment models and insurance models in response to the feedback. These superior risk-assessment models and insurance models can include information about the risk-related behavior, risk-related trends, or forward-looking risk-assessment of selected individuals or selected subsets of the population. These superior risk-assessment models and insurance models can be responsive to data-mining techniques described in related patent applications, described below, hereby incorporated by reference as if fully 10 set forth herein. These superior risk-assessment models can also incorporate known 11 scientific information regarding health risk or disease progression, such as well-12 determined correlations of risk factors and disease incidence or progression from large 13 research studies, or well-known shape of 5-year survival curves for patients having specific types of cancer. 15.

Accordingly, it would also be advantageous to provide a set of techniques for modeling and scoring risk-assessment and a set of insurance products derived therefrom, using dynamic assessment of risk indicators and associated consequences for a population. This advantage is achieved in an embodiment of the invention in which a population (such as a population of medical patients) is assessed both at a selected time and afterward for those risk indicators and for consequences associated therewith. A

WO 00/17799

PCT/US99/22019

client-server system provides dynamic data collection and analysis, dynamic risk assessment in response to that data collection and analysis, and dynamic treatment

options and utilization review for each population member.

Summary of the Invention

The invention provides a set of techniques for modeling and scoring risk-assessment and a set of insurance products derived therefrom. A set of risk indicators (such as medical risk factors for individuals) is determined at a selected time. A population (such as a population of medical patients) is assessed at the selected time and afterward for those risk indicators and for consequences associated therewith. For example, the population can be periodically assessed for correlation between smoking and heart disease, for correlation between alcohol use and heart disease, and for multivariate correlation of a plurality of such indicators and consequences.

In a preferred embodiment, selected population members are each coupled to client devices for determining risk indicators and consequences. For example, where the population is a set of medical patients, the client device can include a local device for asking medical, psychological and life-style questions, and for measurement of medical parameters, for each of those patients. A server device receives data from each client device, and in response thereto, can (1) reassess weights assigned to the risk indicators, (2) determine new significant risk indicators, (3) determine new significant measures for

member or for selected population subsets.

determining risk indicators and consequences, and (4) present treatment options to each.

population member. The server device can perform these tasks in conjunction with an

operator, such as a skilled medical professional, risk-management assessor, or other

expert.

5

The server device can determine, in response to the data from each client device, and possibly in response to other data (such as provided by the expert operator), a measure of risk for each indicated consequence or for a set of such consequences. The server device can provide this measure with regard to each population member, or with regard to population subsets (selected either with regard to the known risk indicators or other indicators). The expert operator can use this measure to determine either (1) an individual course of treatment, (2) a resource utilization review model, (3) a risk-assessment model, or (4) an insurance pricing model, for each individual population

15

16

17

18

19

20

21

22

In a preferred embodiment, information requested by the client device, information determined and presented by the server device, and measurements determined in response thereto, can be adapted dynamically to changing aspects or changing membership of the population, or of an external environment having relevance to the population. For example, medical treatment or risk-assessment models can be dynamically adapted to an aging population or to biomedical advances with regard to detection or treatment of medical conditions for members of that population.

ľ

Brief Description of the Drawings

	3	þ		
	2	,		

Figure 1a shows a block diagram of a system for data collection and interpretation for a population. Figure 1b shows details of the client device 110 shown in figure 1a.

Figure 1c shows devices that may be connected to client device 110. Figure 1d shows details of the data review device.

R

Figure 2 shows a response diagram of consequences to risk indicators, for statistical aggregates of the population, which can be selected in response to dynamic data collection and analysis.

12

13

15

16

Figure 3a shows a process flow diagram of a method for dynamic data collection to be performed by the system; verification of model, updating a model, or creating a new model, and re-evaluation of risk assessment. Figure 3b shows a process flow diagram of the step of dynamic data collection. Figure 3c shows a process flow diagram of the step of verification of the model. Figure 3d shows a process flow diagram of the step of updating the existing model.

19

20

Figure 4a shows a process flow diagram of a method for dynamic data analysis to be performed by the system. Figure 4b shows a process flow diagram for data mining.

.22

•	
Ż	Figure 5 shows a response diagram of consequences to risk indicators, for
3 -	statistical aggregates of the population, with data collected from an individual at different
4 _: .	points of time also plotted.
5	
6.	Figure 6 shows a process flow diagram for a method of providing treatment
7	options and information to each patient based on the data provided to the server.
8.	
 9	Detailed Description of the Preferred Embodiment
0	
1	In the following description, a preferred embodiment of the invention is described
2	with regard to preferred process steps and data structures. Embodiments of the invention
3	can be implemented using general purpose processors or special purpose processors
4	operating under program control, or other circuits, adapted to particular process steps and
5	data structures described herein. Implementation of the process steps and data structures
6	described herein would not require undue experimentation or further invention.
7	
8	Related Applications
9	

Inventions described herein can be used in combination or conjunction with

inventions described in the following patent applications:

21

14

15

18

	of Stephen I Brown, titled
•	Application Serial No. 09/041,809 filed in the name of Stephen J. Brown, titled
	"Phenoscope and Phenobase," assigned to the same assignee, attorney docket
	number RYA-136 and related application serial no. 08/946,341.
	Application Serial No. 07/977,323, filed November 17, 1992 in the name of
	Stephen J. Brown, and issued April 26, 1994 as Patent No. 5,307,263, titled

Stephen J. Brown, and issued April 26, 1994 as Patent No. 5,307,263, titled "Modular Microprocessor Based Health Monitoring System," assigned to the same assignee; and subsequent Continuation-in-Part applications including Application Serial No. 08/481,925 filed June 7, 1995 and Application Serial No. 08/233,397 filed April 26, 1994, and a Continuation-in-Part application filed August 19, 1998, serial number unknown.

Application Serial No. 09/127,404 filed July 31, 1998 in the name of Stephen J. Brown, titled "Modular Microprocessor Based Diagnosed Measurement System for Psychological Conditions", and previous applications of which this is a continuation including Application Serial No. 08/843,495, filed April 16, 1997, which is a continuation of Application Serial No. 08/682,385 filed July 15, 1996, which is a continuation of Application Serial No. 08/479,570 filed June 7, 1995, which is a continuation of Application Serial No. 08/233,674 filed April 26, 1994.

Application Serial No. 08/666,242 filed June 20, 1996, in the name of Stephen
 J. Brown, titled "Health Management Process Control System", assigned to the same assignee, attorney docket number RYA-114.

wo	(100/1	17799

,	Application Serial No. 08/669,613 filed June 24, 1996, in the names of Stephen
2	J. Brown and Erik K. Jensen, titled "On-line Health Education and Feedback
3	System Using Motivational Driver Profile Coding and Automated Content
4	Fulfillment", attorney docket no. RYA-115.
5	• Application Serial No. 08/732,158 filed October 16, 1996, in the name of
. 6	Stephen J. Brown, titled "Multiple Patient Monitoring System for Proactive
7	Health Management", attorney docket no. RYA-116.
8	• Application Serial No. 08/814,293 filed March 10, 1997, in the name of
9	Stephen J. Brown, titled "On-Line Health Education Using Composites of
10	Entertainment and Personalized Health Information", attorney docket no.
11	RYA-119.
12	• Application Serial No. 08/847,009 filed April 30, 1997, in the name of Stephen
<i>13</i>	J. Brown, titled "Monitoring System for Remotely Querying Individuals",
14	attorney docket no. RYA-126.
15	• Application Serial No. 08/975,774 filed in the name of Stephen J. Brown, titled
16	"Multi-User Remote Health Monitoring System", attorney docket no. RYA-
7	.131.
8	and
9	Application Serial No, Express Mail Mailing No. E1027453472US, filed
o .	September 23, 1998, in the name of Stephen J. Brown, titled "Reducing Risk
	\cdot

	17799

PCT/US99/22019

l	- 4

Using Behavioral and Financial Rewards," assigned to the same assignee, attorney docket number HHN-004.

3

These applications are hereby incorporated by reference as if fully set forth herein.

5

System for Data Collection

7

Figure 1a shows a block diagram of a system for data collection and interpretation for a population.

Referring to figure 1a, a system 100 includes a client device 110, a server device
11 120 including a program memory 122 and database of patient information 121, and a data
12 review element 130. These devices are connected via a communication channel, such as
13 a communication network as in known in the art and more fully described in the
14 Phenoscope and Phenobase patent (U.S. 09/041,809) and related patent application serial
15 no. 08/946,341 and other patents and patent applications previously incorporated by
16 reference.

Referring to figure 1b, the client device 110 is disposed locally to a patient 111,

and includes an output element 112 for presenting information to the patient 111, and an

input element 113 for entering information from the patient 111. As used herein,

"locally" refers to a logical relationship to the patient 111, and does not have any

necessary implication with regard to actual physical position. In a preferred embodiment,

WO 00/17799 PCT/US99/22019

the client device 110 is relatively small or compact, and can be disposed on a night table

or otherwise near the patient 111.

3

The output element 112 includes a display screen 114, on which questions and suggested answers can be displayed for the patient 111, so as to facilitate information entry, or on which instructions can be displayed for the patient 111, so as to instruct the

patient 111. The output element 112 can also include a speaker 115, so as to present

8 information in conjunction with or in alternative to the display screen 114. The output

9 element 112 can also include a bell or other sound element, or a bright light 119 or a flag,

so as to alert the patient 111 that the client device 110 has questions or information for

11 the patient 111.

12

13

14

15

10

The input element 113 includes a plurality of buttons 116A-D for entering information, preferably such as described in the patent applications referenced and incorporated by reference above.

16

17

18

. 19

20

21

22

The input element 113 can also include one or more data ports 117A-D for entering information from other devices. Referring to figure 1c, such other devices 118 can include a medical measurement device, such as a blood glucose meter or a blood pressure monitor. Such other devices 118 can include a dispensing device for medication.

2	Such other devices 118 can also include a general purpose or special purpose
. 3	client workstation, such as a personal computer or a hand-held digital calendar.
. 4	
5	The server device 120 is disposed logically remotely from the patient 111, and
6	includes a database 121 of information about the patient 111 and about other patients in
7	related population thereof. As used herein, "remotely" refers to a logical relationship to
8	the patient 111, and does not have any necessary implication with regard to actual
9	physical position.
10	
11	The server 120 and patient profile database 121 are preferably accessible by mean
12	of a standard network connection such as a world wide web connection. Server 120 and
13	database 121 may comprise single stand-alone computers or multiple computers
] 4	distributed throughout a network.
15	
16	Referring to figure 1a and figure 1d, the data review element 130 is disposed
17	logically remotely from the patient 111, and includes an interface 131 disposed for use by
18	an operator 132. The operator 132 can comprise medical personnel, a device operated by
19	medical personnel, or a similar device, capable of interacting with the interface 131 so as
20	to receive information from the data review element 130 and possibly to enter

information into the data review element 130. Information entered into the data review

2	client device 110.
3	
	The data review element 130 is preferably a personal computer, remote terminal,
· ·	
.5	web TV unit, Palm Pilot unit, interactive voice response system, or any other
· 6	communication technique. The data review element functions as a remote interface for
7	entering in server 120 or client device 110 messages and queries to be communicated to
8	the individuals.
9 .	Other and further information regarding the system 100 is shown in the following
10	pending patent applications and in other patent applications referenced above:
11	
12	• Application Serial No. 09/041/809, filed in the name of Stephen J. Brown
13	titled "Phenoscope and Phenobase," assigned to the same assignee, attorne
14	docket number RYA-136 and related application serial no. 08/946,341.
15	and
16	Application Serial No, Express Mail Mailing No. EI027453472US
17	filed September 23, 1998, in the name of Stephen J. Brown, titled "Reducin
-18	Risk Using Behavioral and Financial Rewards," assigned to the same assignee
19	attorney docket number HHN-004.
20	
21	These applications are hereby incorporated by reference as if fully set forth herein
22	

Aggregate Responses to Risk Indicators

ż

Figure 2 shows a response diagram 200a of consequences to risk indicators, for statistical aggregates of the population, which can be selected in response to dynamic

- data collection and analysis. It is to be noted that figure 2 shows curves that are collapsed
- 6 to 2-dimensions, in a preferred embodiment the curves are N-dimensional, with N>2.

7

- 8 A diagram 200a includes a first axis X 201 and a second axis Y 202. The diagram
- 9 shows a first response curve R0 210 showing a normal trajectory for vital function and
- 10 life expectancy of an individual or subpopulation of the population. The first axis X 201
- 11 indicates a relative time, as measured toward a right side of the diagram. The scale of the
- 12 first axis X 201 is a relative time whose initial left hand point may be undetermined. As
- 13 to a first response curve R0 210, the second axis Y 202 represents a measure of vital
- 14 function and life expectancy.

15

- 16 A diagram 200a also shows a second response curve S0 220 showing a normal
- 17 trajectory for a measure of expected medical expense or risk for an individual or
- 18 subpopulation of the population. The first axis X 201 indicates a relative time as for a
- 19 first response curve R0 210. As to a second response curve S0 220, the second axis Y
- 20 202 shows increasing expense or risk as measured toward the top of the diagram.

In the first response curve R0 210, the normal trajectory for vital function and life

- expectancy for a typical individual in the population shows that as time progresses,
- vitality and life expectancy are expected to decrease. This general concept is known in
- the art of actuaries. It is to be noted that the shape shown by the first response curve R0
- 210 is an example shape; for instance, it is known that for certain curable cancers, risk
- increases, then levels off after a certain length of time such as a 5-year survival rate, then
- later in life risk increases due to other causes.

11

12

13

14

15

16

17

18

19

20

21

The first response curve R0 210 includes a number of points with error bars 211 about the response curve R0 210. All of the points 211 are at an identical value, V0, of the second axis Y 202, with identical error bars. Any one of the points represents a single measurement of vitality taken for an individual. Given any single measurement of vitality, it is difficult to determine where along the second axis X 201, that is, where along the trajectory the individual is. Of particular interest is how close to a rapid decline in vitality or increase in risk the individual is. The points 211 show the several places along the curve where the individual might be placed, based on this single measurement of vitality. Because the response curve R0 210 is slowly varying through much of the time, that is, the values of vitality and life expectancy clustering in a selected region of the second axis Y 202, shown by the bracket 203, and due to margins of error in both the measurement as well as the response curve, there are several positions along the curve where an individual with a specific measurement might be; these several positions are shown by points 211.

By contrast, if measurements are taken for an individual at more than one point in time, greater information is present, and in particular trends may be discerned which yield more information about where on the curve an individual is. This ability to discern trends is greater when curves in N-dimensions are considered. For instance, an individual whose excess weight has slowly climbed in conjunction with slowly increasing cholesterol, blood pressure, stress levels and family medical history would be placed in a greater risk category although the individual measures of, for instance, cholesterol, might be within a normal range. .10 Similarly, in the second response curve S0 220, the normal trajectory for expected medical expense and risk for that typical individual shows that as time progresses, 12 expected medical expense and risk are expected to increase. This general concept is also known in the art of actuaries. It is to be noted that the shape shown by the second response curve R0 220 is an example shape; for instance, upon diagnosis of a disease the expense may climb, but if the patient is cured the expense will level off. Similarly, the second response curve S0 220 includes a number of points 221 on the response curve S0 220, showing possible places that an individual in the population

17

18

19

20

21

22

13

15

16

with measurement of expense or risk, with value E0, might be. Because most of the values of response curve S0 220 cluster in a selected region of the second axis Y 202 shown by the bracket 204, it is difficult to know where along curve S0 220 an individual

- r with measurement E0 should be placed. This is due to both possible error in
- 2 measurement of E0 as well as uncertainty in the exact "true" position and shape of curve
- 3 SO 220. As for curve RO 210, measurements of expense or risk taken over time will yield
- 4 useful information about where on the curve SO 220 an individual is.
- When subsets of the population are selected in response to specific risk factors, the
- 7 statistical aggregates of the population can differ substantially from the aggregate
- 8 response curves R0 210 and S0 220 for the entire population. The diagram 200a shows
- 9 response curves R1a 212 and R1b 213 showing a normal life trajectory for vital function
- and life expectancy of an "average" individual in the population, depending on whether
- 11 that individual is associated with a selected risk factor a. As with regard to the aggregate
- 12 for the entire population, it is difficult to determine from a specific single measurement
- just where on either response curve R1a 212 or R1b 213 the individual should be
- assessed. Depending on whether the value of α is known for an individual, it may also be
- difficult to know whether the individual should be placed on response curve R1a 212 or
- 16 R1b 213. Measurements of several risk indicators taken over time may yield information
- on whether a specific individual should be placed in category R1a 213 or the higher risk
- 18 category R1b 212. The general concept of using time-dependent information to
- 19 determine risk along is also illustrated in Figure 5.

or

	The client device 110 determines information from which the server device 120 or
	z the data review element 130 can analyze the time varying nature of data. The server
	device 120 or the data review element 130 can therefore determine both of the following:
	•
٠.,	• (1) just where on either response curve R1a 212 or R1b 213 the individual
•	should be assessed; and
. ,	
8	• (2) whether the individual should be assessed on the response curve R1a 212 or
9	
10	
. 11	It is to be noted that the above analysis has been condensed to 2-dimensions for
12	convenience in presentation, with a single measurement along a single X-axis or Y-axis.
13	In a preferred embodiment, a measurement would have many attributes, i.e. the model
14	would have N-dimensions, and more sophisticated techniques for analyzing trends and
15	achieving objectives are used.
16	
17	If the data for the population is not known for all individuals in the population or
18	subpopulation of interest, the server device 120 transmits a new set of information-
19	gathering instructions (such as questions and suggested answers) to the client device 110,
20	so as to measure that information individually for each patient 111.
21	

١.	Dynamic	Mode	eling.and	Risk	Evaluation
----	---------	------	-----------	------	------------

Figure 3a shows a process flow diagram 300a for a method with steps of
dynamically collecting information 310, choosing to verify or update the model or to
create new model 320, verifying 350 or updating 330 the risk assessment model or
creating a new model 340, deciding whether to re-evaluate risk 360 and re-evaluating risk
based on updated information and current model 370.

7

Dynamic Data Collection for Population

Figure 3b shows a process flow diagram 300b of a method for dynamic data collection to be performed by the system. This data collection may be done periodically or aperiodically, upon a triggering event or decision by the expert operator. The population or subpopulation from which to collect data is selected 380. The selection criteria may be based on preset values or may be set by the expert operator. The set of risk indicators or other information to be collected is selected 382, based either on preset values or decision by the expert operator. The individuals in the subpopulation of interest are queried 384 as to the information of interest and the database is updated 386. The pre-query steps need not be done in the order indicated.

18

Verification of Existing Model and Update of Model

2Ò

Figure 3c shows a process flow diagram 300c of a method by which the updated
data can be analyzed to determine whether the existing model is consistent with the

updated data; that is, to verify that the data conforms to the model within acceptable

variation or error. This is accomplished by putting the updated data into categories 390,

determining the updating measures of life vitality or costs 392, determining the values

1 predicted by the model 394, comparing the updated measures of life vitality or costs

5 against those predicted by the model 396, and determining whether the comparison is

6 acceptable 397. If the predicted value is within an acceptable distance from the updated

values based on well known measures such as statistical error, then the model need not be

8 adjusted. The expert operator may also visually determine whether the updated data and

existing model show an acceptable relationship to each other.

10

15

16

17

18

19

20

Figure 3d shows a process flow diagram 300d of a method for updating the existing risk model in response to updated information. By updating, it is meant that no new risk indicators are added, and no new external constraints on the model are added. The risk model to be adjusted may be for the aggregate population or for various subpopulations. The updated information for the subpopulation is categorized 398 according to profile information into one or more existing categories. The subpopulation is categorized according to one or more existing measure of life vitality or medical expense. Statistical analyses as described below or in other patents or patent applications previously incorporated by reference or as known in the art of statistics are applied to determine updated values for model parameters such as weights to give each factor 399.

2]

,	was asked of the client but was not previous	sly known to b	e a significant	predictor, or it
2	may be a new factor that is generated by co	mbining other	pieces of data.	Figure 4b is a
,	process flow diagram of the above steps.	• • • • • • • • • • • • • • • • • • • •		

In addition to data mined from the database, in creating a new model, scientific information well known in the literature may supplement the data. For instance, scientific information regarding certain well studied correlations be considered such as known correlations of time since quitting smoking and various health conditions, known information regarding the shape of life expectancy curves for certain types of cancer patients, or recent information regarding efficacy of new forms of treatment for diseases such as recent significant improvements in treatment of AIDS.

Statistical analyses are known in the art of statistics, and include correlation analyses, multivariate regressions, constrained multivariate regressions, or variance analyses, may also be run on the data to reveal statistical relationships among the various information or measures of life vitality or medical expense in order to improve the predictive power of a model, although in a preferred embodiment data mining is done as presented in the preceding paragraphs.

Modeling and Scoring Risk Assessment, Insurance Pricing

÷ 1	Modeling risk is performed by assigning risk to individual in response to risk
2	factors identified for that individual, and such modeling may be done for the population
3	or for a subpopulation. There are many techniques for modeling, such as linearly risk
4	scoring by assigning a number to each risk factor and adding up each number to
5	determine a total risk score, non-linearly assessing risk by combining risk factors non-
6	linearly to determine risk which may be achieved by neural network techniques which a
. 7	known in the art of neural networks, or other techniques.
8	
9	Figure 5 shows a diagram 500 including a first axis X 502 and a second axis Y
10.	503 and a response curve R0 501, similar to that shown in figure 2. It shows several
11	measurements of vitality with error bars 511 of an individual taken at several different
12	points in time. Each measurement of vitality is taken at a later time from left to right.
13	Information about the time varying nature of the measurements, or the trends, can
14	improve the ability to predict future vitality, including imminent sharp declines in
15	vitality, as can been seen by visually examining the data over time or by using
16	sophisticated statistical techniques to examine the data and trends in the data over N-

19

17

dimensions.

Insurance pricing may be achieved from advantages in risk assessment. It is known in the art of actuarial analysis to assign price in response to risk.

21

<i>i</i> .			· 21		•	
	•			• •		
?	Providing treatment opti	ions and inf	ormation to ea	ch populatio	n member	

Figure 6 is a process flow diagram 600 showing a method for providing treatment options and information to each member based on the information provided. Upon

receiving information about the patient from the client 610, the server or expert operator

may identify a risk group 620 and identify an appropriate medical protocol 630, the

server may present one or more responses to the patient 640, including treatment options,

advice or merely health information that would be useful to the patient, and the client

device may be configured to use an appropriate medical protocol in interacting with the

patient 650. It is known in the art of medicine that membership in a risk group may 10

indicate appropriate treatment. This may be done from an automated, preset set of

responses to individual queries made to the patient, on an aggregate of preset responses to

queries, or by an expert operator. 13

Alternative Embodiments

12

14

16

21

Although preferred embodiments are disclosed herein, many variations are possible which remain within the concept, scope, and spirit of the invention, and these 18 variations would become clear to those skilled in the art after perusal of this application. 19

28 Claims

•	
3	1. A method for assessing risk for selected individuals in a population, said method
4	including steps for
5	determining, at a first time, a first set of risk indicators for said
6	selected individuals;
7	collecting, at a second time after said first time, information about
8	said selected individuals;
9.	determining, at said second time, an additional risk indicator not in
10.	said first set, in response to said information;
11	assessing risk for said selected individuals in response to said
12	additional risk indicators.
13	
14,	2. A method for assessing risk for selected individuals in a population, said method
15	including steps for
16	determining, at a first time, a set of risk indicators for said selected
17	individuals;
j8	collecting, at a second time after said first time, information about
19	said selected individuals;
20	adjusting, at said second time, at least one of said risk indicators in
21	response to said information;
	· · · · · · · · · · · · · · · · · · ·

20

	29 assessing risk for said selected individuals in response to said
. 2	adjusted risk indicators.
4	3. A method as in claim 2, wherein said risk indicators include genetic risk
5	indicators, medical risk indicators, environmental risk indicators, or behavioral risk
. 6	
. 7	
8	4. A method as in claim 2 wherein said stone for a new and stone
9	to be an ordinal 2, wherein said steps for confecting include steps for
	collecting, at said second time, information for said selected individuals about a set o
10	consequences associated with said risk indicators.
"	
12	5. A method as in claim 2, including steps for determining a statistical
. 13 .	measure of relation between at least one said risk indicator and said information about
14	said selected individuals.
15	
16	6. A method as in claim 2, including steps for determining a statistical
17	measure of relation between at least two said risk indicators and said information about
18	said selected individuals.
19	
20	7. A method as in claim 2, wherein said steps for collecting include steps for
21	providing a client device for at least one of said selected individuals;

NAC 00/1/2200		•	PCT/US99/2201
₩O 00/17799			1 (1/03)7/2201

,		applying a measurement device to said one selected individual at
2	• 4	said client device;
3		coupling said client device to a server device; and
4		transmitting a result of said steps for applying to said server device.
5	÷	
6	8.	A method as in claim 2, wherein said steps for collecting include steps for
7		providing a client device for at least one of said selected individuals;
8	·· .	displaying questions at said client device; and
9		receiving answers to said questions from said at least one selected
0		individual;
1		
2	9.	A method as in claim 8, wherein said steps for displaying include steps for
3	•	receiving said questions from a server device coupled to said client
4		device;
s [.]		timing said steps for displaying in response to a signal from said
6 _		server device; and
7	•	transmitting said answers to said server device.
8		
9	10.	A risk-assessment model, said model including
Ó .		a set of risk indicators for selected individuals in a population;
1		a first set of values associated, at a first time, with each
2		corresponding risk indicator;

. 1		a set of information associated, at a second time after said first time,
2		with said selected individuals;
.3	•	a second set of values associated, at said second time, with each said
4		corresponding risk indicator, said second set of values being determined in
5		response to said set of information;
6		a risk-assessment, determined in response to said second set of
7	·. '.	values, for said selected individuals.
8		
9	11.	A financial product including
10		a set of risk indicators for selected individuals in a population;
11	. : .	a first set of values associated, at a first time, with each
12	• • •	corresponding risk indicator;
13	•	a set of information associated, at a second time after said first time,
14		with said selected individuals;
15		a second set of values associated, at said second time, with each said
16		corresponding risk indicator, said second set of values being determined in
17 ·		response to said set of information;
18		a pricing value, determined in response to said second set of values,
19		for said selected individuals.
20		
21	12.	A financial product as in claim 11, wherein said pricing value is an insurance premium





Fig. 1A






Figure 1c




Figure 1d






Figure 24




Egune 3a



300b

Figure 36



300c

Figure 3c Venification Categorize updated

Data According

to existing

nuclei categories


398

Optimize existing model parameters to fit data subject to existing model Constraints

399

3001

Update Existing Risk Model Figure 31






Figure 46

WO 00/17799 12 / 13 PCT/US99/22019




Fig. 5

Server Receives Information From Client

610

Server or Expert.
Operator Identifies
Risk Group ASSESSA

620

gener or Expert
Operator Identifies
Appropriate Medical
Protocol

630

Response according to
Appropriate Medical
Protocol or as set
by Expert Operator
is communicated to
client

640

Client May Be
PRICEAL Configured to
Use that Medical
Robocol in Francisco

650

France

600

INTERNATIONAL SEARCH REPORT

PCT/US 99/22019

	·		PU1/05 9	PC1/02 AA\5501A		
A. CLASS	FICATION OF SUBJECT MATTER G06F17/60				•	
v ,			٠,			
According t	to International Patent Classification (IPC) or to both national cla	essification and IPC				
. FIELDS	SEARCHED			<u> </u>		
	ocumentation searched (classification system followed by class	silication symbols)				
IPC 7	606F				•	
)ocumenta	tion searched other than minimum documentation to the extent	that such documents are inch	uded in the fields	searched		
			٠.			
	tata base consulted during the international search (name of da	to have and where provided	against toring trac	<u> </u>		
:lectronic c	1919 Dase Consumed during the memberonia search (maine or or	na pase and, where precinces	' attach intina dad	u ,	٠.	
	ENTS CONSIDERED TO BE RELEVANT					
	CRation of document, with indication, where appropriate, of the	ho relevant nace anne		Relevant to claim f	No.	
category *	CERTION IN COCCUMENT, MAINTENANCE, MAINTENANCE, OF D	To room a passages	,		<u>.</u>	
	50 0 700 000 A (MINOUS 2011 0505	7 (44 94000)		1-12		
(·	EP 0 700 009 A (MINGUIJON PERE	Z SALVADUK)		1-12		
	6 March 1996 (1996-03-06) column 1, line 55 -column 2, l	ine 54		1		
		inc 51	· //			
(PALFREY T R ET AL: "Repeated	insurance		1-4,		
	contracts and learning"		•	10-12	• . •	
	RAND JOURNAL OF ECONOMICS, AUT	UMN 1985,				
	USA,		•		. `	
	vol. 16, no. 3, pages 356-367	•				
	XP000878736 ISSN: 0741-6261					
	page 356, line 17 -page 357, 1	ine 6				
	,				٠.	
		-/				
				·		
	•					
			•			
			•		•	
				L <u> </u>		
X Furth	er documents are listed in the continuation of box C.	X Patent family n	nembers are listed	in annex.		
Special cel	egories of cited documents :	<u> </u>	 		•	
•		"I" later document publi or priority date and	ished after the lints not in conflict with	rnational filing date the application but		
" docume: conside	nt defining the general state of the lart which is not ered to be of particular relevance	cited to understand invention	the principle or th	eory underlying the		
earlier d tiling da	ocument but published on or after the international	"X" document of particul		laimed invention		
documer	nt which may throw doubts on priority claim(s) or	cannot be consider involve an inventive	ed novel or cannot slep when the do	criment is laken alone .		
which is citation	s cited to establish the publication date of another or other special reason (as specified)	"Y" document of particul	ar relevance; the d	taimed invention ventive step when the		
	nt reterring to an oral disclosure, use, exhibition or	document is combi	ned with one or ma	ore other such docu- us to a person skilled		
ather in documen	of published prior to the international filing date but	in the art.				
later the	an the priority date claimed	"8" document member o				
ate of the a	ctual completion of the international search	Date of mailing of the	ne international se	arch report		
		05 /00 /00				
15	February 2000	25/02/20				
eme and m	ailing address of the ISA	Authorized officer				
	European Patent Office, P.B. 5818 Patentiaan 2	· ·		•. •		
	Nt. – 2280 HV Rijswijk Tel. (+31–70) 340–2040, Tr., 31 651 epo nl,	Pedersen	ı. N	,	1	
	Fax: (+31-70) 340-3016	1 16061361	• • • • • • • • • • • • • • • • • • • •			

INTERNATIONAL SEARCH REPORT

PCT/US 99/22019

		PC1/05 99	722019	
(Continu	NION) DOCUMENTS CONSIDERED TO BE RELEVANT			•
ategory *	Citation of document, with indication, where appropriate, of the relevant passages		Relevant to claim No.	
	BAEHRING T U ET AL: "Using the World Wide		1-10	
`	Web—a new approach to risk identification	•		
	of diabetes mellitus"			
	INTERNATIONAL JOURNAL OF MEDICAL			
	INFORMATICS, IR, ELSEVIER SCIENTIFIC			
	PUBLISHERS, SHANNON,			
	vol. 46, no. 1,	•		
	1 August 1997 (1997-08-01), pages 31-39,			
	XP004085528	:		
	ISSN: 1386-5056	•		
	page 34, column 2, line 1 -page 35, column			
· ·	1. line 12			
	1, Fille 12		ŀ	
P, X.	MONTANI S ET AL: "Protocol-based		1-10	
「 ,^	reasoning in diabetic patient management"			
٠.,	INTERNATIONAL JOURNAL OF MEDICAL		1 .	
	INTERNATIONAL GOODRAL OF MEDICAL INFORMATICS, IR, ELSEVIER SCIENTIFIC	•		
ļ	PUBLISHERS SHANNON.	•	٠ .	
,	vol. 53, no. 1, January 1999 (1999-01),			
	pages 61-77, XP004158051			`.
	ISSN: 1386-5056]	
	page 63, column 1, line 19 -page 64,			·
	column 1. line 11			
١ .	CLEMONS E K ET AL: "Information		1-12	
`	technology and information asymmetry: the			
	future of private individual health			
- 1	insurance"	• *	* *	٠.
	PROCEEDINGS OF THE THIRTIETH HAWAII			
1	INTERNATIONAL CONFERENCE ON SYSTEM			
	SCIENCES (CAT. NO.97TB100234), PROCEEDINGS			
1	OF THE THIRTIETH HAWAII INTERNATIONAL		·	
	CONFERENCE ON SYSTEM SCIENCES, WAILEA, HI,			
	USA, 7-10 JAN. 1997, pages 240-248 vol. 3,		·	
l	XP002130528			
l	1997, Los Alamitos, CA, USA, IEEE Comput.			
	Soc. Press, USA ISBN: 0-8186-7743-0			
i	page 240, column 2, line 40 -page 241,	:	٠.	
	column 1, line 27		İ	
ı	page 242, column 1, line 9 - line 27			
· .	6B 2 231 420 A (AGENCY MANAGEMENT SERVICES		1-12	•
`	INC) 14 November 1990 (1990–11–14)	•		
ı	page 8, line 18 -page 9, line 5			
ł	hade at time to hade at time of	•		
!				٠.
ľ	•			
ŀ	. •			•
l l				
ı				
- 1				
[
- 1		,		
į				
	•	•		
1	•			

INTERNATIONAL SEARCH REPORT

-information on patent family member

Intel 2011 Applics. No PCT/US 99/22019

Patent document cited in search report		Publication date	Patent lamily member(s)		Publication date	
-	EP 0700009	À	06-03-1996	ES	2108613 A	16-12-1997
	GB 2231420	A	14-11-1990	NONE		

--- DCT//CA/710 (pagers family annes) (July 1997)